arXiv:hep-th/9906196v3 20 Jan 2000

KUNSAN-TP-99-6

Absorption of dilaton s-wave in type 0B string theory

Jin Young Kim*
Department of Physics, Kunsan National University, Kunsan, Chonbuk 573-701, Korea

Abstract

We find the absorption probability of dilaton field in type 0B string theory. Since the background solutions are of the form $AdS_5 \times S^5$ on both regions, we use the semiclassical formalism adopted in type IIB theory to find the absorption cross section. The background tachyon field solution was used as a reference to relate the solutions of the two regions. We also consider the possible corrections to absorption probability and the $\ln(\ln z)$ form of the correction is expected as in the calculation of the confinement solution.

PACS number(s): 04.70.Bw, 11.27.+d

Typeset using REVTEX

---

*Electronic address: jykim@ks.kunsan.ac.kr

I. INTRODUCTION

Maldecena's conjecture that $\mathcal{N} = 4$ super Yang-Mills theory is dual to type IIB supergravity has opened a new horizon to the study of large N QCD [1]. Another possibility to apply this technique to non supersymmetric gauge theories was proposed by Polyakov [2]. He suggested that one can study non-SUSY gauge theories using branes in the context of type 0 string theories. The type 0 string theories are defined on the world sheet of type II by performing a non-chiral GSO projection [3]. As a consequence type 0 theories have world sheet supersymmetry but no space-time supersymmetry. The resulting sectors of the theories are, in the notation of [4], $(NS-, NS-), (NS+, NS+)$ and a doubled set of Ramond-Ramond fields. If the doubled R-R is $(R+, R-), (R-, R+)$, the theory is type 0A and if $(R+, R+), (R-, R-)$, the theory is 0B. These non-SUSY string theories are also part of the web of dualities and can be described as limits of M-theory. It is known that the type 0A corresponds to M-theory on $\mathbf{S}^1/(-1)^{F_s} \cdot S$, where S denotes a half-shift along the circle, and type 0B corresponds to M-theory on $\mathbf{T}^2/(-1)^{F_s} \cdot S$, where S is along the circle of the torus [5].

The low energy fields of the theory are tachyon, the bosonic field of type II and two copies of R-R fields. Accordingly they have open string descendants which are crucial for the D-brane construction. Following the idea of [2], Klebanov and Tseytlin [6] sought the possibility that the tachyonic instability can be cured by the R-R fields. Using the near-horizon threebrane geometry they constructed an $SU(N)$ gauge theory with six adjoint scalars and studied its behavior in the point of AdS/CFT correspondence. The tachyon indeed condensed with its effective mass squared being shifted to a positive value. The analysis showed that the tachyon is constant at large radial distance and the geometry is of the form $AdS_5 \times S^5$ with vanishing coupling, which can be interpreted as a UV fixed point. The condensation breaks the conformal invariance which agrees with the fact that the dual gauge theory is not conformal. The asymptotic solution for weak coupling was found by Minahan [7] and revealed that the logarithmic flow as expected from the asymptotic freedom of the dual gauge theory. Also the solution has the right sign of the β-function in agreement with field theory and there exists another fixed point at infinite coupling(IR fixed point), which is also characterized by $AdS_5 \times S^5$ Einstein frame metric. It is argued that there is an RG trajectory that extends from UV fixed point to the IR fixed point [8,9]. The generic static solution to type 0 gravity equations of motions are confining in the IR and have logarithmic scaling in the UV.

The construction of non-SUSY branes suggests many directions of research. And it seems that many of the results which were obtained in type II brane configurations can be copied in the type 0 case. Since the metric solutions are asymptotically $AdS_5 \times S^5$ for both regions, the study of scattering process can be one of the examples. Scattering amplitude or absorption cross section involving the tachyon field cannot be addressed from the previous work on type II theories [10,11]. However one can use the same technique as in type II theories [12] to evaluate various scattering with D-branes in type 0 theories. The purpose of this paper is to seek this possibility. Specifically, we will consider the semiclassical low energy scattering of dilaton under the static background solution of type 0B string theory and find the absorption cross section.

The organization of the paper is as follows. In Section II we review the type 0B theory and its background solutions of both regions and set up some preliminaries for our calculation. In section III we find the leading absorption cross section in the low-frequency limit. We use the background tachyon field as a reference to relate the solutions of both regions. In Section IV we consider the higher order corrections to the absorption probability. Two kinds of orrections are possible, one from the background metric and the other from the string coupling through the tachyon field. Section V is devoted to conclusions.

II. TYPE 0B BACKGROUND SPACETIME

The type 0 model has a closed string tachyon, no fermions and a doubled set of R-R fields, and thus a doubled set of D-branes [13]. With the doubling of R-R fields, the self-dual constraint on the five-form field is relaxed and one can have D3 branes that are electric instead of dyonic. We consider the low energy world volume action of N coincident electric D3 branes. We start from the action for the type 0B theory [8]

$$\begin{aligned} S_0 = & -\frac{1}{2\kappa_{10}^2}\int d^{10}x\sqrt{-g}\Big[R-\frac{1}{2}\nabla_n\Phi\nabla^n\Phi \\ & -\frac{1}{4}(\nabla_nT\nabla^nT+m^2e^{\frac{1}{2}\Phi}T^2)-\frac{1}{4\cdot 5!}f(T)F_{n_1\ldots n_5}F^{n_1\ldots n_5}+\cdots\Big], \end{aligned} \tag{1}$$

where g_{mn} is the Einstein-frame metric, $m^2=-\frac{2}{\alpha'}$, and the tachyon-R-R field coupling function is

$$f(T)=1+T+\frac{1}{2}T^2. \tag{2}$$

The equations of motion from this effective action are

$$\nabla^2\Phi=\frac{1}{8}m^2e^{\frac{1}{2}\Phi}T^2, \tag{3}$$

$$\begin{aligned} R_{mn}-\frac{1}{2}g_{mn}R = & \frac{1}{4}\nabla_mT\nabla_nT-\frac{1}{8}g_{mn}[(\nabla T)^2+m^2e^{\frac{1}{2}\Phi}T^2]+\frac{1}{2}\nabla_m\Phi\nabla_n\Phi \\ & -\frac{1}{4}g_{mn}(\nabla\Phi)^2+\frac{1}{4\cdot 4!}f(T)(F_{mklpq}F_n{}^{klpq}-\frac{1}{10}g_{mn}F_{sklpq}F^{sklpq}), \end{aligned} \tag{4}$$

$$(-\nabla^2+m^2e^{\frac{1}{2}\Phi})T+\frac{1}{2\cdot 5!}f'(T)F_{sklpq}F^{sklpq}=0, \tag{5}$$

$$\nabla_m[f(T)F^{mnkpq}]=0. \tag{6}$$

If one parametrize the 10-d Einstein-frame metric as in [6]

$$ds_E^2=e^{\frac{1}{2}\xi-5\eta}d\rho^2+e^{-\frac{1}{2}\xi}(-dt^2+dx_idx_i)+e^{\frac{1}{2}\xi-\eta}d\Omega_5^2, \tag{7}$$

where ρ is the radial direction transverse to the 3-brane ($i=1,2,3$), then the radial effective action corresponding to (3) – (6) becomes

$$S = \int d\rho \Big[\frac{1}{2}\dot{\Phi}^2 + \frac{1}{2}\dot{\xi}^2 - 5\dot{\eta}^2 + \frac{1}{4}\dot{T}^2 - V(\Phi, \xi, \eta, T) \Big], \tag{8}$$

$$V = \frac{1}{2\alpha'} T^2 e^{\frac{1}{2}\Phi + \frac{1}{2}\xi - 5\eta} + 20 e^{-4\eta} - Q^2 f^{-1}(T) e^{-2\xi}. \tag{9}$$

Here the constant Q is the R-R charge and dot means the derivative with respect to ρ. The resulting set of variational equations,

$$\ddot{\Phi} + \frac{1}{4\alpha'} T^2 e^{\frac{1}{2}\Phi + \frac{1}{2}\xi - 5\eta} = 0, \tag{10}$$

$$\ddot{\xi} + \frac{1}{4\alpha'} T^2 e^{\frac{1}{2}\Phi + \frac{1}{2}\xi - 5\eta} + 2Q^2 f^{-1}(T) e^{-2\xi} = 0, \tag{11}$$

$$\ddot{\eta} + 8e^{-4\eta} + \frac{1}{4\alpha'} T^2 e^{\frac{1}{2}\Phi + \frac{1}{2}\xi - 5\eta} = 0, \tag{12}$$

$$\ddot{T} + \frac{2}{\alpha'} T e^{\frac{1}{2}\Phi + \frac{1}{2}\xi - 5\eta} + 2Q^2 \frac{f'(T)}{f^2(T)} e^{-2\xi} = 0, \tag{13}$$

should be supplemented by the 'zero-energy' constraint

$$\frac{1}{2}\dot{\Phi}^2 + \frac{1}{2}\dot{\xi}^2 - 5\dot{\eta}^2 + \frac{1}{4}\dot{T}^2 + V(\Phi, \xi, \eta, T) = 0, \tag{14}$$

which can be used instead of one of the second-order equations.

In the Einstein frame the dilaton decouples from the R-R terms ($|F_5|^2$) and the tachyon mass term plays the role of the source term. It is easy to analyze the case where the $|F_5|^2$ is large in (5). Assuming that the tachyon is localized near the extremum of $f(T)$, i.e. $T = -1$, the asymptotic solution in the UV($\rho \equiv e^{-y} \ll 1,\ y \gg 1$) region is given by [7,8]

$$\bar{T} = -1 + \frac{8}{y} + \frac{4}{y^2}(39 \ln y - 20) + O(\frac{\ln^2 y}{y^3}), \tag{15}$$

$$\bar{\Phi} = \ln(2^{15} Q^{-1}) - 2 \ln y + \frac{39}{y} \ln y + O(\frac{\ln y}{y^2}), \tag{16}$$

$$\bar{\xi} = \ln(2Q) - y + \frac{1}{y} + \frac{1}{2y^2}(39 \ln y - 104) + O(\frac{\ln^2 y}{y^3}), \tag{17}$$

$$\bar{\eta} = \ln 2 - \frac{1}{2} y + \frac{1}{y} + \frac{1}{2y^2}(39 \ln y - 38) + O(\frac{\ln^2 y}{y^3}). \tag{18}$$

The 10-d Einstein frame metric can be written as

$$
\begin{aligned}
ds_E^2 = R_0^2 \Big[& (1 - \frac{9}{2y} - \frac{351}{4y^2}\ln y + \cdots)(\frac{1}{4}dy)^2 \\
& + (1 - \frac{1}{2y} - \frac{39}{4y^2}\ln y + \cdots)\frac{e^{\frac{1}{2}y}}{2R_0^4} dx^\mu dx^\mu + (1 - \frac{1}{2y} - \frac{39}{4y^2}\ln y + \cdots) d\Omega_5^2 \Big],
\end{aligned} \tag{19}
$$

where

$$R_0^2 = 2^{-1/2} Q^{1/2}.$$

Note that with $y = 4\ln u$, one can show that the metric is of the form of $AdS_5 \times S^5$ at the leading order,

$$ds_E^2 = R_0^2 (\frac{du^2}{u^2} + \frac{u^2}{2R_0^4} dx^\mu dx^\mu + d\Omega_5^2). \tag{20}$$

The corrections cause the effective radius of AdS_5 to become smaller than that of S^5. One can find the asymptotic freedom from the large u behavior of the leading effective gravity solution. It is an important question whether it survives the full string theoretic treatment. It has been argued [8] that the solution does survive due to the special structure related to the approximate conformal invariance. The crucial fact is that the Einstein metric is asymptotic to $AdS_5 \times S^5$. This geometry is conformal to flat space, so that the Weyl tensor vanishes in the large u limit. Furthermore, both Φ and T vary slowly for large u.

One interesting feature of the tachyon RG trajectory is that T starts increasing from its critical value $T = -1$ from condition $f'(T) = 0$. The precise form of the trajectory for finite ρ is not known analytically, but the qualitative feature of the RG equation was analyzed by Klebanov and Tseytlin [8,14]. Since Φ, ξ and η have negative second derivatives (see equations (10) –(12)), each of these fields may reach a maximum at some value of ρ. If, for Φ, this happens at finite ρ, then the coupling decreases far in the infrared. As a result, a different possibility can be realized: $\dot{\Phi}$ is positive for all ρ, asymptotically vanishing as $\rho \to \infty$. They constructed the asymptotic form of such trajectory. The point is that, as $\rho \to \infty$, T approaches zero so that, as in the UV region, $T^2 e^{\frac{1}{2}\Phi}$ becomes small. For this reason the limiting Einstein-frame metric is again $AdS_5 \times S^5$. Thus, the theory flows to a conformally invariant point at infinite coupling. The explicit form of the asymptotic large ρ solution (IR solution) is

$$\bar{T} = -\frac{16}{y} - \frac{8}{y^2}(9\ln y - 3) + O(\frac{\ln^2 y}{y^3}), \tag{21}$$

$$\bar{\Phi} = -\frac{1}{2}\ln(2Q^2) + 2\ln y - \frac{9}{y}\ln y + O(\frac{\ln y}{y^2}), \tag{22}$$

$$\bar{\xi} = \frac{1}{2}\ln(2Q^2) + y + \frac{9}{y} + \frac{9}{2y^2}(9\ln y - \frac{20}{9}) + O(\frac{\ln^2 y}{y^3}), \tag{23}$$

$$\bar{\eta} = \ln 2 + \frac{1}{2}y + \frac{1}{y} + \frac{1}{2y^2}(9\ln y - 2) + O(\frac{\ln^2 y}{y^3}), \tag{24}$$

where y is related to ρ by $\rho \equiv e^y (\gg 1,\ y \gg 1)$. The 10-d Einstein frame metric in the IR region can be written as

$$ds_E^2 = R_\infty^2 \Big[(1 - \frac{1}{2y} - \frac{9}{4y^2}\ln y + \cdots)(\frac{1}{4}dy)^2 \\ +(1 - \frac{9}{2y} - \frac{81}{4y^2}\ln y + \cdots)\frac{e^{-\frac{1}{2}y}}{2R_\infty^4}dx^\mu dx^\mu + (1 + \frac{7}{2y} + \frac{63}{4y^2}\ln y + \cdots)d\Omega_5^2\Big], \tag{25}$$

where

$$R_\infty^2 = 2^{-3/4}Q^{1/2}.$$

The metric starts again as $AdS_5 \times S^5$ ($y = -4\ln u$) at $y = \infty$ and becomes a negative curvature 10-d space at smaller y (bigger u). As in the large u region, the corrections cause the effective radius of AdS_5 to become smaller than that of S^5.

The structure of the asymptotic UV and IR solutions is similar, suggesting that they can be smoothly connected into the full interpolating solution. For example, the tachyon starts at $T = -1$ at $\rho = 0$, and grows according to (15), then enters an oscillating regime and finally relaxes to zero according to (21). Note that the coordinates corresponding to the two regions are related by $y \to -y$.

III. SEMICLASSICAL ABSORPTION OF DILATON

If we consider the solution found in section II as a static background solution, one can think the low energy scattering of string fields. Since the metric solutions are asymptotically $AdS_5 \times S^5$ in both regions, we can use the semiclassical formalism of type II theories [12]. For simplicity of calculation we will consider scattering of low energy dilaton field in the type 0B string backgrounds. To study the scattering problem, we introduce the time-dependent perturbed field of dilaton

$$\Phi = \bar{\Phi}(y) + \phi(y,t). \tag{26}$$

We look for the solutions to the equation of motion for a mode of frequency ω of the dilaton $\phi(y,t) = \phi(y)e^{i\omega t}$. Then the linearized equation of motion governing the perturbation of the dilaton is

$$\frac{1}{\sqrt{-g}}\partial_y\Big(\sqrt{-g}g^{yy}\partial_y\phi(y)\Big) - \omega^2 g^{tt}\phi(y) + \frac{1}{8\alpha'}e^{\frac{\bar{\Phi}}{2}}\bar{T}^2\phi(y) = 0. \tag{27}$$

We will consider the solution in the UV region first. To find the leading order solution in the UV region, we insert the background metric (19) in the limit $y \to \infty$, which is $AdS_5 \times S^5$, into the equation

$$16\frac{d^2\phi}{dy^2} + 16\frac{d\phi}{dy} + 2\omega^2 R_0^4 e^{-\frac{1}{2}y}\phi + \frac{R_0^2}{8\alpha'}e^{\frac{\bar{\Phi}}{2}}\bar{T}^2\phi = 0, \tag{28}$$

where the background values of $\bar{T}$ and $\bar{\Phi}$ are given by (15) and (16). This equation becomes in terms of $u = \exp(y/4)$

$$\frac{d^2\phi}{du^2} + 5\frac{d\phi}{du} + \frac{2\omega^2 R_0^4}{u^2}\phi + \frac{R_0^2}{8\alpha'} e^{\frac{\bar{\Phi}}{2}} \bar{T}^2 \phi = 0. \tag{29}$$

We introduce new dimensionless variable

$$z \equiv \frac{\sqrt{2}\omega R_0^2}{u} = \sqrt{2}\omega R_0^2 e^{-\frac{y}{4}} \tag{30}$$

and putting $\phi = \psi(z)/u^2$, we have

$$z^2 \frac{d^2\psi}{dz^2} + z\frac{d\psi}{dz} + (z^2 - 4)\psi + \frac{R_0^2}{8\alpha'} e^{\frac{\bar{\Phi}}{2}} \bar{T}^2 \psi = 0. \tag{31}$$

In the UV region where the string coupling is weak, the last term is negligible and the leading order solution is

$$\psi(z) = H_2^{(2)}(z), \tag{32}$$

$$\phi(z) = \left(\frac{z}{\sqrt{2}\omega R_0^2}\right)^2 H_2^{(2)}(z), \tag{33}$$

where we choose the solution pure infalling at the horizon.

Now we consider the solution in the IR region. Inserting the background metric (25) in the limit $y \to \infty$ into the equation, we have

$$16\frac{d^2\phi}{dy^2} - 16\frac{d\phi}{dy} + 2\omega^2 R_\infty^4 e^{\frac{1}{2}y}\phi + \frac{R_\infty^2}{8\alpha'} e^{\frac{\bar{\Phi}}{2}} \bar{T}^2 \phi = 0. \tag{34}$$

Here the background values of $\bar{T}$ and $\bar{\Phi}$ are given by the solutions in the IR region (21) and (22). Introducing new variable in the IR region as $v = \exp(y/4)$, the equation becomes

$$\frac{d^2\phi}{dv^2} - 3\frac{d\phi}{dv} + \frac{2\omega^2 R_\infty^4}{v^2}\phi + \frac{R_\infty^2}{8\alpha'} e^{\frac{\bar{\Phi}}{2}} \bar{T}^2 \phi = 0. \tag{35}$$

Note that u and v are related by inverse because y's in the UV and IR regions are related by $y \to -y$. In terms of new dimensionless variable

$$\sigma \equiv \sqrt{2}\omega R_\infty^2 v = \sqrt{2}\omega R_\infty^2 e^{\frac{y}{4}} \tag{36}$$

and with $\phi = v^2 f(\sigma)$, we have

$$\sigma^2 \frac{d^2 f}{d\sigma^2} + \sigma\frac{df}{d\sigma} + (\sigma^2 - 4)f + \frac{R_\infty^2}{8\alpha'} e^{\frac{\bar{\Phi}}{2}} \bar{T}^2 f = 0. \tag{37}$$

In the IR region, we can also neglect the last term since $\bar{T} \to 0$ in the leading order and the solution is given by the Bessel functions

$$f(\sigma) = \alpha J_2(\sigma) + \beta J_{-2}(\sigma), \tag{38}$$

where the first term corresponds to the normalizable mode and the second to nonnormalizable mode. We choose $\beta = 0$ for the solution to be finite as $\sigma \to \infty$, then

$$\phi(\sigma) = \alpha\Big(\frac{\sigma}{\sqrt{2}\omega R_\infty^2}\Big)^2 J_2(\sigma). \tag{39}$$

To relate the solution in the UV region and IR region, we adopt the matching scheme similar to the cases of D3-brane and M5-branes [10] and D1-D5 brane system [11]. For both cases the equation of motion for the minimal scalar has a self-dual point defined by the radius of the effective anti-de Sitter space. In our case we have to be careful in choosing the matching point because the effective radii of the anti-de Sitter space for both regions are different because of the tachyon coupling. However, the tachyonic coupling is always combined with the background dilaton $\bar{\Phi}$, i.e. the coupling is of the form of $e^{\bar{\Phi}/2}\bar{T}^2$. And it seems that the functional behavior of the coupling terms for both UV and IR region are the same. To see this, we calculate the coupling terms up $O(1/y)$. Substituting the UV background solution (15) and (16) into the coupling term, we have

$$R_0^2 e^{\frac{\bar{\Phi}}{2}} \bar{T}^2 = 2^{-\frac{1}{2}} Q^{\frac{1}{2}} \cdot 2^{\frac{15}{2}} Q^{-\frac{1}{2}} \frac{1}{y} \cdot 1^2 = \frac{2^7}{y}. \tag{40}$$

Similarly, from (21) and (22), we have

$$R_\infty^2 e^{\frac{\bar{\Phi}}{2}} \bar{T}^2 = 2^{-\frac{3}{4}} Q^{\frac{1}{2}} \cdot 2^{-\frac{1}{4}} Q^{-\frac{1}{2}} y \cdot \frac{16}{y}^2 = \frac{2^7}{y}. \tag{41}$$

Up to the order of $O(1/y)$, the functional form of the perturbed terms are the same. And this may be true for higher orders too. This implies that the functional forms of the correction will be the same. Thus if we extend the UV solution toward the IR region and IR solution toward the UV region, they can be matched at some point of y. But we don't know the exact position of y. So we choose the matching point from the point where the background tachyon fields have the same values at the leading order. Equating (15) and (21) in the leading order, $-1 + 8/y = -16/y$, we have $y = 24$ as the matching point. In terms ρ, this corresponds that we match the asymptotic solution of the UV region, which is valid for very small ρ, at $\rho = e^{-24}$ and that of the IR region, valid for very large ρ, at $\rho = e^{24}$. Matching the amplitude of ϕ at the leading order

$$\Big(\frac{z}{\sqrt{2}\omega R_0^2}\Big)^2 H_2^{(2)}(z)\Big|_{z=\sqrt{2}\omega R_0^2 e^{-6}} = \alpha\Big(\frac{\sigma}{\sqrt{2}\omega R_\infty^2}\Big)^2 J_2(\sigma)\Big|_{\sigma=\sqrt{2}\omega R_\infty^2 e^{-6}}, \tag{42}$$

one finds the relative coefficient relating the solutions of two regions.

$$\alpha = \frac{8i}{\pi}\frac{e^{24}}{\omega^4 R_0^4 R_\infty^4} \propto \frac{1}{\omega^4 Q^2}. \tag{43}$$

The asymptotic form of the infalling wave function in the UV region is

$$\phi(z) = \Big(\frac{z}{\sqrt{2}\omega R_0^2}\Big)^2 H_2^{(2)}(z) \simeq \frac{1}{(\sqrt{2}\omega R_0^2)^2}\sqrt{\frac{2}{\pi}} z^{\frac{3}{2}} \exp\{-i(z - \frac{3}{4}\pi)\}. \tag{44}$$

From this the invariant flux, defined by

$$F_{\rho\to 0} = \frac{1}{2i}(\phi^* \sqrt{-g} g^{yy}\partial_y \phi - \phi\sqrt{-g} g^{yy}\partial_y \phi^*)\Big|_{y=\infty}, \tag{45}$$

is given by

$$F_{\rho\to 0} = \frac{1}{2\pi}. \tag{46}$$

Since the ingoing part of the wavefunction in the IR region is

$$\phi(\sigma) = \alpha\Big(\frac{\sigma}{\sqrt{2}\omega R_\infty^2}\Big)^2 J_2(\sigma) \simeq \frac{1}{(\sqrt{2}\omega R_\infty^2)^2}\sqrt{\frac{1}{2\pi}}\sigma^{\frac{3}{2}}\exp\{i(\sigma - \frac{3}{4}\pi)\}, \tag{47}$$

the ingoing flux at infinity is given by

$$F_{\rho\to\infty} = \frac{|\alpha|^2}{8\pi} = \frac{8e^{48}}{\pi^3\omega^8 R_0^8 R_\infty^8}. \tag{48}$$

The absorption probability is given by the ratio of the flux in the UV region to the ingoing flux in the IR region and we find

$$P = \frac{F_{\rho\to 0}}{F_{\rho\to\infty}} = \frac{\pi^2\omega^8 R_0^8 R_\infty^8}{16e^{48}} \propto \omega^8 Q^4. \tag{49}$$

The s-wave absorption cross-section is given by [12]

$$\sigma^s_{abs} = \frac{(2\pi)^{d-1}}{\omega^{d-1}\Omega_{d-1}}P\Big|_{d=6} = \frac{\pi^4}{e^{48}}R_0^8 R_\infty^8 \omega^3 \propto \omega^3 Q^4. \tag{50}$$

Note that the cross section has an ω^3 dependence, similar to the type II case. If one chooses different position of y as the matching point, the numerical coefficient can be changed but the ω and Q dependence will be the same.

IV. HIGHER ORDER CORRECTIONS

In this section we will consider the effect of higher order corrections of the dilaton absorption cross-section. The dominant corrections to absorption cross-section arise from the matching about the point where the tachyon field have the same value. In the case of D3-brane and M5-brane of type IIB theory the correction to the minimally coupled scalar equation is done by considering the higher order terms of the background metric. However, in our case where the dilaton is coupled to the tachyon through the string coupling α', we have to consider the correction by this coupling term too. For simplicity of the calculation we consider only the leading terms and subleading corrections will be straightforward.

In the UV region, inserting

$$\sqrt{-g} = \frac{R_0^2 e^y}{16}(1 - \frac{9}{2y} + \cdots),\ g^{yy} = \frac{16}{R_0^2}(1 + \frac{9}{2y} + \cdots),\ g^{tt} = -2R_0^2 e^{-\frac{1}{2}y}(1 + \frac{1}{2y} + \cdots)$$

and

$$\bar{T}^2 = 1 - \frac{16}{y} + \cdots,\quad e^{\frac{\bar{\Phi}}{2}} = \frac{2^7}{R_0^2}(\frac{1}{y} + \cdots)$$

into (27), we have

$$16\frac{d^2\phi}{dy^2}+16\frac{d\phi}{dy}+2\omega^2R_0^4e^{-\frac{1}{2}y}(1+\frac{1}{2y}+\ldots)\phi+\frac{16}{\alpha'}(\frac{1}{y}+\cdots)\phi=0. \tag{51}$$

Substituting $y = 4\ln u$ and $\phi = \psi/u^2$, this can be written as

$$\frac{d^2\psi}{du^2}+\frac{d\psi}{du}+(\frac{2\omega^2R_0^4}{u^2}-4)\psi=\Big\{\frac{2\omega^2R_0^2}{u^2}(\frac{4}{y}+\cdots)-\frac{16}{\alpha'}(\frac{1}{y}+\cdots)\Big\}\psi. \tag{52}$$

With the dimensionless variable defined earlier $z = \frac{\sqrt{2}\omega R_0^2}{u} = \sqrt{2}\omega R_0^2 e^{-\frac{y}{4}}$, we get

$$z^2\frac{d^2\psi}{dz^2}+z\frac{d\psi}{dz}+(z^2-4)\psi=\Big\{\frac{z^2}{\ln\frac{\sqrt{2}\omega R_0^2}{z}}(1+\cdots)-\frac{4}{\alpha'}\frac{1}{\ln(\frac{\sqrt{2}\omega R_0^2}{z})}(1+\cdots)\Big\}\psi. \tag{53}$$

In the low energy scattering case the terms on the right hand side act as small perturbations. The first term on the right hand side is due to metric correction while the second is from the tachyon coupling. We look for the perturbative solution of the form [10,11,15]

$$\psi(z)=\psi_0(z)+\psi_1(z)+\cdots, \tag{54}$$

where the zeroth order solution is

$$\psi_0(z)=H_2^{(2)}(z). \tag{55}$$

ψ_1 satisfies the inhomogeneous equation

$$z^2\frac{d^2\psi_1}{dz^2}+z\frac{d\psi_1}{dz}+(z^2-4)\psi_1=\Big\{\frac{z^2}{\ln\frac{\sqrt{2}\omega R_0^2}{z}}(1+\cdots)-\frac{4}{\alpha'}\frac{1}{\ln\frac{\sqrt{2}\omega R_0^2}{z}}(1+\cdots)\Big\}\psi_0. \tag{56}$$

The solution to this equation is

$$\psi_1(z)=\frac{\pi}{2}\int^z dx\frac{1}{x}\Big(\frac{-x^2}{\ln\frac{\sqrt{2}\omega R_0^2}{x}}+\frac{4}{\alpha'}\frac{1}{\ln\frac{\sqrt{2}\omega R_0^2}{x}}\Big)H_2^{(2)}(x)\{J_2(x)Y_2(z)-J_2(z)Y_2(x)\}. \tag{57}$$

Substituting the series form of Bessel functions, and retaining the terms necessary for given order, one can find the correction in series form.

We repeat the same procedure in the IR region to find the higher order solution of the form

$$f(\sigma)=f_0(\sigma)+f_1(\sigma)+\cdots, \tag{58}$$

where $f_0(\sigma)$ is the zeroth order solution given by $f_0(\sigma) = \alpha J_2(\sigma)$. The inhomogeneous equation which f_1 should satisfy is

$$\sigma^2\frac{d^2f_1}{d\sigma^2}+\sigma\frac{df_1}{d\sigma}+(\sigma^2-4)f_1=-\Big\{\frac{\sigma^2}{\ln\frac{\sigma}{\sqrt{2}\omega R_\infty^2}}(1+\cdots)+\frac{4}{\alpha'}\frac{1}{\ln\frac{\sigma}{\sqrt{2}\omega R_\infty^2}}(1+\cdots)\Big\}f_0, \tag{59}$$

and the solution is

$$f_1(\sigma) = \alpha\frac{\pi}{2}\int^{\sigma} dx\frac{1}{x}\Big\{\frac{x^2}{\ln\frac{x}{\sqrt{2}\omega R_\infty^2}}(1+\cdots) + \frac{4}{\alpha'}\frac{1}{\ln\frac{x}{\sqrt{2}\omega R_0^2}}(1+\cdots)\Big\}J_2(x)\{J_2(x)Y_2(\sigma) - J_2(\sigma)Y_2(x)\}. \tag{60}$$

Matching the two solutions at the matching point

$$\Big(\frac{z}{\sqrt{2}\omega R_0^2}\Big)^2(\psi_0(z)+\psi_1(z))\Big|_{z=\sqrt{2}\omega R_0^2 e^{-6}} = \alpha\Big(\frac{\sigma}{\sqrt{2}\omega R_\infty^2}\Big)^2 f_0(\sigma) + f_1(\sigma)\Big|_{\sigma=\sqrt{2}\omega R_\infty^2 e^{-6}}, \tag{61}$$

one can find the correction of the relative coefficient α.

If we substitute the series expansion of Bessel functions, it is expected the that the leading correction is of the form $\ln(\ln z)$ from the integral structure

$$\int^z \frac{dx}{x}\frac{x^n}{\ln x} = z^n \ln(\ln z) + \cdots.$$

However, if we retain only the leading terms of the the series expansion of Bessel functions

$$J_2(x) = \frac{x^2}{8},\quad Y_2(x) = -\frac{4}{\pi x^2},\quad H_2^{(2)}(x) = \frac{4i}{\pi z^2}, \tag{62}$$

the correction terms from this type cancels out. The correction of the relative coefficient α is from the subleading terms i.e. next order terms of Bessel functions and $\cdots$ of (57) and (60). This is a good consistency check for the background solutions.

V. DISCUSSION

We considered the low energy scattering of dilaton field in the type 0B theory. Using the fact that the asymptotic structure of the background solutions on both regions is $AdS_5 \times S^5$, we calculated semiclassically the absorption probability. Since the tachyonic coupling term is small on both regions, the calculation is similar to the case of minimally coupled scalar of type II theory. A crucial role in our formalism was played by the background tachyon field. We used it as a reference to find the matching point. We also considered the possible corrections of absorption probability and the $\ln(\ln z)$ form of the correction [7,8] is expected as in the calculation of the confinement solution.

In the present paper, we perturbed only the dilaton field from the background for simplicity of calculation. The right way to analyze the scattering is to perturb all fields in the theory and solve the full coupled equations, which seems very difficult to solve. We considered the low energy world volume action of N coincident electric D3 branes. We expect that the same formalism can be applied to self-dual 3-branes [14] because the asymptotic structure of the background also has $AdS_5 \times S^5$ structure. Though we considered only the case of $d = 10$ critical case, it seems interesting to study the case of non critical type 0 theory in lower dimensions [16].

ACKNOWLEDGEMENT

I would like to thank S.P.Kim, Y.S.Myung and H.W.Lee for useful discussions. This work was supported by the Korean Physical Society (1999).

REFERENCES

[1] J. Maldacena, Adv. Theor. Math. Phys. **2**, 231(1998), hep-th/9711200; S.S. Gubser, I.R. Klebanov and A.M. Polyakov, Phys. Lett. **B428**, 105(1998), hep-th/9802109; E. Witten, Adv. Theor. Math. Phys. **2**, 253(1998), hep-th/9802150.

[2] A.M. Polyakov, Int. J. Mod. Phys. Phys. **A14**, 645(1999), hep-th/9809057.

[3] L. Dixon and J. Harvey, Nucl. Phys. **B274**, 93(1986); N. Seiberg and E. Witten, Nucl. Phys. **B276**, 272(1986).

[4] J. Polchinski, *String Theory*, Vol. 2 (Cambridge University Press, 1998).

[5] O. Bergman and M.R. Gaberdiel, hep-th/9906055.

[6] I.R. Klebanov and A.A. Tseytlin, Nucl. Phys. **B546**, 155(1999), hep-th/9811035.

[7] J.A Minahan, JHEP **9901**, 020(1999), hep-th/9811156.

[8] I.R. Klebanov and A.A. Tseytlin, Nucl. Phys. **B547**, 143(1999), hep-th/9812089.

[9] J.A. Minahan, JHEP **9904**, 007(1999), hep-th/9902074.

[10] S.S. Gubser, A. Hashimoto, I.R. Klebanov, and M. Krasnitz, Nucl. Phys. **B526**, 393(1998), hep-th/9803023.

[11] M. Taylor-Robinson, hep-th/9806132; A. Rajaraman, Phys. Lett. **B437**, 69(1998), hep-th/9803082; S. Mathur and A. Matusis, hepth/9805064.

[12] I. Klebanov, Nucl. Phys. **B496**, 231(1997), hep-th/9702076.

[13] O. Bergman and M.R. Gaberdiel, Nucl. Phys. **B499**, 183(1997), hep-th/9701137.

[14] I.R. Klebanov and A.A. Tseytlin, JHEP **9903**, 015(1999), hep-th/9901101.

[15] J.Y. Kim, H.W. Lee and Y.S. Myung, hep-th/9812016.

[16] G. Ferretti and D. Martelli, hep-th/9811208; A. Armoni, E. Fuchs and J. Sonnenschein, hep-th/9903090; G. Ferretti, J. Kalkkinen and D. Martelli, hep-th/9904013.